



## White Paper 23-08

### Genetic Associations with Traits in 23andMe Customers

---

*Authors:*

Saniya Fayzullina

Robin P. Smith

Nicholas Furlotte

Youna Hu

David Hinds

Joyce Y. Tung

*Created:* December 4, 2014

*Revised:* February 25, 2015

June 3, 2015

(see end of document for summary of major changes)

### Summary

**Purpose:** This white paper serves as a companion to 23andMe customer reports, offering more details and scientific support for the reported findings. We hope that the paper may also serve the scientific community to elaborate molecular pathways and contribute to the understanding of basic human biology and disease.

**Methods:** Customers submitted saliva for DNA extraction and testing using the 23andMe Personal Genome Service® (PGS®). The 23andMe PGS® uses a BeadChip

array to probe over half a million genetic variants distributed across the entire human genome. Trait data were gathered through questions presented via the 23andMe website. Anonymized trait and genotype data were then combined to conduct a GWAS. Associations meeting the genome-wide significance criterion for each trait ( $p < 5 \times 10^{-8}$ ) are incorporated into customer reports.

Findings: We report the following trait associations:

- *Cilantro Aversion:* Two genetic loci are significantly associated with aversion to cilantro. Both loci are located in a cluster of olfactory receptor genes that have been linked with the ability to detect aldehyde odorants.
- *Sweet vs. Salty/Savory Taste Preference:* Two genetic loci that are significantly associated with the preference for sweet-tasting foods. These loci are located within the *FGF21* and *FTO* genes, both of which are associated with metabolism and body mass regulation.
- *Sensitivity to the Sound of Chewing (Misophonia):* One genetic locus that is significantly associated with the sensitivity to chewing sounds. This locus is located near the *TENM2* gene that plays a role in brain development.

Conclusions: We will continue to perform GWAS on a variety of traits and use data reported in this white paper to support customer reports. We will continue adding new trait associations to this white paper as more 23andMe GWAS results become available.

## Introduction

The 23andMe Personal Genome Service® (PGS®) uses DNA extracted from customer saliva to give customers access to their genetic information. Customers are invited to answer questions about various traits and health conditions. Their anonymized survey responses and genetic data are combined to perform a genome-wide association study (GWAS).

By combining our large consented customer research database, web-based surveys, and custom GWAS tools, we routinely identify genetic associations for a variety of human traits. Some of these analyses have been used to replicate associations reported in the literature and to publish novel associations in peer reviewed journals [1–13].

This white paper details genetic associations that we have identified with the participation of 23andMe customers. It is intended to support customer reports that provide interpretations of personal genetic results. We will update this white paper over time as we add more customer reports.

## Methods

Single-phenotype GWAS were conducted as described previously [6]. Relevant details and changes are described briefly below.

### Study Participants

Study participants were 23andMe customers who had been genotyped as part of the 23andMe PGS®, consented to research, and voluntarily responded to web-based surveys. The study protocol and consent form were approved by the external Association for the Accreditation of Human Research Protection Programs-accredited Institutional Review Board, Ethical and Independent Review Services.

All study participants were required to have > 97% European ancestry, as determined by analysis of local ancestry [14]. The reference population data for ancestry analysis were derived from public datasets (the Human Genome Diversity Project, HapMap, and 1000 Genomes) and from 23andMe customers who have reported having four grandparents from the same country. At present, the database has the highest power to detect associations in cohorts of European ancestry.

We also required that participants were unrelated (sharing a smaller percent of the genome than the minimum expected for first cousins), as determined by a segmental identity-by-descent estimation algorithm [15].

### Sample collection, DNA extraction, and genotyping

Participants provided their saliva using a specialized collection device, and mailed their samples to a CLIA-certified laboratory contracted by 23andMe. DNA extraction and 23andMe BeadChip genotyping were performed by the contracted laboratory [1,6]. All samples were genotyped on one of four versions of 23andMe BeadChips (produced by Illumina) that probe over half a million single nucleotide polymorphisms (SNPs) distributed across the entire human genome.

### Trait data collection

Each GWAS presented in this paper is based on the response to one or more questions presented via the 23andMe website. Each question is designed to collect data about a specific trait or a set of related traits. Questions are typically multiple choice. The 23andMe self-report data gathered through such questions is statistically robust and replicable, as shown by a number of collaborator publications [3,6,9,11,16–18].

### GWAS

Single-phenotype GWAS were conducted as previously described [6], except that no phasing or SNP imputation were done. Associations between genotyped SNPs and traits were estimated using logistic regression assuming additive allelic effects for binary traits (to calculate an odds ratio (OR)) and linear regression for quantitative traits (to calculate a regression coefficient  $\beta$ ). The analysis was adjusted for age, sex, and the top five principal components to account for population structure.  $P$  values were computed using a likelihood ratio test. SNPs were excluded from the GWAS analysis if any of the following criteria were met: minor allele frequency  $< 0.1\%$ , Hardy-Weinberg equilibrium test [19]  $p < 10^{-20}$  in people of European ancestry, evidence of laboratory batch effects, or genotype call rate  $< 95\%$ . SNPs were also excluded if they had large allele frequency discrepancies compared with European 1000 Genomes reference data, identified by computing a 2x2 table of allele counts of European 1000 Genomes samples and 2000 randomly sampled 23andMe customers with European ancestry, and identifying SNPs with a chi squared  $p < 10^{-15}$ . After GWAS analysis,  $p$  values were adjusted using genomic control (a factor of 1.092 for cilantro preference data, 1.227 for sweet preference data, and 1.103 for misophonia data) to correct for statistical inflation [20]. SNPs with an association  $p < 5 \times 10^{-8}$  were considered statistically significant.

### Trait selection for reporting

For customer reports, we gave higher priority to associations that had larger effect sizes, identified novel phenotype associations, or had strong evidence for underlying biological mechanisms.

### Data presentation

Association  $p$  values for all genotyped SNPs across all chromosomes are illustrated in a Manhattan plot. Compelling associations will have clusters of significantly associated SNPs at one or more chromosomal loci, with a few SNPs with highly statistically significant  $p$  values at each locus. Data quality was further assessed in the Q-Q plot, which plots observed quantiles of GWAS  $p$  values against expected quantiles of  $p$  values under a null distribution. To identify index SNPs, all SNPs with association  $p < 10^{-5}$  were grouped into intervals separated by gaps of at least 250kb. The SNP with the lowest statistically significant  $p$  value in each interval was considered to be the index SNP for that region, and reported in Table 1. SNP positions are specified according to the NCBI Genome Reference Consortium human genome build 37 (GRCh37).

## Associations identified by GWAS

### Cilantro taste aversion

The *Coriandrum sativum* plant is commonly known as cilantro or coriander. It is used as an ingredient in various cuisines around the world, with historical references even found in ancient Roman texts [21–23]. However, some people describe cilantro taste as unpleasant and “soapy”, and the prevalence of cilantro aversion appears to vary with ancestry [24]. The key chemical components responsible for cilantro aroma are thought to be aldehydes: unsaturated aldehydes are described as smelling “fruity” and (E)-2-alkenals are described as “soapy” [25–27]. These and many other odorants are detected by specialized odorant receptors, products of a family of about 400 functional human genes [28].

Genetic differences in odorant receptors have been shown to affect how we perceive tastes and smells [1,29–35]. To learn more about potential genetic associations, we assessed participants’ perception of cilantro taste using a multiple-choice question with 3 answer options (in parentheses):

*“Do you like the taste of fresh (not dried) cilantro?” (Yes/No/I’m not sure)*

A total of 56,360 participants (74%) answered “Yes” and 19,731 participants (26%) answered “No”. Participants who answered “I’m not sure” were excluded.

We identified 2 genomic loci that passed criteria for genome-wide association significance (summarized in Figure 1, detailed data in Table 1). Index SNPs for these loci, rs2741762 and rs3930459, lie in non-coding regions within a cluster of 56 olfactory receptor genes in chromosomal region 11p15.4 [36]. The A allele of rs2741762 ( $p = 2.1 \times 10^{-14}$ , OR = 1.1) and the C allele of rs3930459 ( $p = 4.2 \times 10^{-13}$ , OR = 1.1) are associated with higher odds of disliking the taste of cilantro.

We previously published a GWAS on cilantro preference using a smaller cohort of 23andMe customers. In that study, we identified a SNP (rs72921001) within 100kb of the OR6A2 olfactory receptor gene associated with perception of a “soapy taste” in cilantro [37]. The data described here represents a refinement of the earlier study methodology with a larger study cohort. The new GWAS identified SNPs within 10kb of the OR10A2, OR10A4, OR10A6, and OR10A3 genes and within 100kb of the OR6A2 gene (noted in the previous study). *OR6A2* is orthologous to the rat *I7* receptor gene [38–42]. The *I7* receptor detects a range of n-aliphatic aldehyde odorants, including aldehyde components of cilantro aroma [27,36,39].

### Sweet versus salty/savory taste preference

Human taste perception depends in part on taste receptors in buds on the tongue [43–45]. Sweet taste preference has approximately 50% heritability [46,47], meaning that genetic and non-genetic factors play an equal role. Studies in children and adults have shown that early childhood experiences and long-term exposures to dietary sugar or salt in adulthood influence one’s perception and preference for sweeter or saltier foods [48–50]. Cultural differences in taste preferences are likely due to differences in the way sugar and salt are used in each cultural group’s cuisine

[51–53]. In addition, there is evidence for interplay between taste preference, food intake, and metabolism, all of which have genetic components [54–58].

To learn more about potential genetic associations with taste preferences, we assessed participants' sweet versus salty/savory food preference using a multiple-choice question with 4 answer options (in parentheses):

*“When you're in the mood for a snack, what kind of snack do you usually reach for?”  
(Sweet / Salty or savory / Both / Neither)*

A total of 54,901 participants (46%) answered “Sweet” and 64,049 participants (54%) answered “Salty or savory”. Participants who answered “Both” or “Neither” were excluded.

We identified 29 genomic loci that passed criteria for genome-wide association significance (summarized in Figure 2). From these we chose two index SNPs with the lowest  $p$  values to be incorporated into customer reports (detailed data in Table 1). The index SNP for the first locus, rs838133, lies within an intron of the gene *FGF21* in chromosomal region 19q13.33. The index SNP for the second locus, rs1421085, lies within an intron of the gene *FTO* in chromosomal region 16q12.2. The A allele of rs838133 ( $p = 1.9 \times 10^{-49}$ , OR = 1.1) and the C allele of rs1421085 ( $p = 3.6 \times 10^{-23}$ , OR = 1.1) are associated with higher odds of preferring sweet tasting foods.

The *FGF21* gene encodes a hormone that regulates glucose and lipid metabolism. [59–62]. In obese individuals, blood levels of FGF21 are elevated and are predictive of developing diabetes [63,64]. Because of its function as a metabolism regulator, FGF21 has been proposed as a therapeutic target for metabolic diseases [65,66].

The *FTO* locus we report was previously found to be associated with obesity [67,68]. This locus contains long-range elements that regulate the expression of the neighboring homeobox gene *IRX3* [69,70]. In turn, *IRX3* is thought to play an important role in the regulation of body mass and metabolism [69].

Our findings are consistent with those of Chu et al., who identified an association between the same *FTO* and *FGF21* loci (including SNP rs838133) and dietary macronutrient intake [71].

### **Sensitivity to the sound of chewing**

“Misophonia” is a very recently coined term to describe a dislike of specific sounds [72]. The offending sounds include the sound of other people chewing or breathing (most common), finger or foot tapping, and footsteps [73]. Misophonia is different from other types of sound sensitivity (such as hypersensitivity to loud sounds) because it has a strong emotional component and the specificity for certain human sounds [74]. Two studies have linked misophonia with a higher frequency of obsessive compulsive personality disorder [73,75]. However, little research has been published on misophonia, and the underlying biology is not known.

To learn more about potential genetic associations with misophonia, we focused on the most common offensive stimulus: the sound of chewing. We assessed participants’ aversion to the sound of chewing using a multiple-choice question with 3 answer options (in parentheses):


“Does the sound of other people chewing fill you with rage?” (Yes/No/Not Sure)

A total of 17,606 participants (22%) answered “Yes” and 63,001 participants (78%) answered “No”. Participants who answered “Not Sure” were excluded.

We identified one genomic locus that passed criteria for genome-wide association significance (summarized in Figure 3, detailed data in Table 1). The index SNP for the significant locus, rs2937573, lies near the gene *TENM2* in chromosomal region 5q34. The G allele of rs2937573 ( $p = 2.0 \times 10^{-39}$ , OR = 1.2) is associated with higher odds of being sensitive to the sound of chewing.

The *TENM2* gene encodes a protein that is involved in the formation of neural connections during brain development [76–78]. One researcher has proposed that misophonia “results from enhanced functional connections between the auditory and the limbic and autonomic nervous systems” [79]. The involvement of *TENM2* supports this theory; however, further investigation is needed to understand the biology of misophonia.


**Figure 1. Cilantro aversion GWAS results**



Manhattan plot of GWAS results for the cilantro taste trait. Log-scaled  $p$  values (y-axis) are plotted against genomic position on chromosomes 1-22 and X (x-axis). Red squares represent SNPs with  $p < 5 \times 10^{-8}$  ( $-\log p$  of  $\sim 7.3$ ), green and blue squares represent SNPs with  $p > 5 \times 10^{-8}$ . The horizontal grey line represents the  $p = 5 \times 10^{-8}$  cut-off. Labels next to the most significant index SNP locations show the names of the nearest genes.




**Figure 2. Sweet vs. salty taste preference GWAS results**



Manhattan plot of GWAS results for the sweet taste preference trait. (See legend for Figure 1.)

**Figure 3. Sensitivity to Chewing Sounds GWAS results**



Manhattan plot of GWAS results for the chewing sound sensitivity trait. (See legend for Figure 1.)



**Table 1. Significant Associations Reported to Customers**

Trait	SNP	Region	Position	Alleles (effect/no effect)	EAF	OR (95% CI)	p value	Gene context
<b>Cilantro aversion</b>	rs2741762	11p15.4	6892866	A/G	0.62	1.102 (1.075 - 1.130)	2.1x10 <sup>-14</sup>	<i>OR10A2</i> [-] <i>-OR10A4</i>
	rs3930459	11p15.4	7953958	C/T	0.41	1.095 (1.069 - 1.122)	4.2x10 <sup>-13</sup>	<i>OR10A6</i> -[-] <i>-OR10A3</i>
<b>Sweet preference</b>	rs838133	19q13.33	49259529	A/G	0.44	1.147 (1.136 - 1.157)	1.9x10 <sup>-49</sup>	[ <i>FGF21</i> ]
	rs1421085	16q12.2	53800954	C/T	0.41	1.096 (1.076 - 1.117)	3.6x10 <sup>-23</sup>	[ <i>FTO</i> ]
<b>Sensitivity to chewing sounds</b>	rs2937573	5q34	166471198	G/A	0.43	1.189 (1.158 - 1.220)	2.0x10 <sup>-39</sup>	[ ]--- <i>TENM2</i>

Index SNPs for regions with association  $p < 5 \times 10^{-8}$  are reported. Region – cytogenetic band position; Position – SNP position according to the NCBI Genome Reference Consortium human genome build 37 (GRCh37); Alleles – effect/no effect on genomic reference strand; EAF – effect allele frequency across all study participants; OR – odds ratio for the high-risk allele; CI – 95% confidence interval; Gene context – gene(s) spanning or flanking (< 1Mb away from) the index SNP; brackets indicate the position of the SNP, and dashes indicate distance to a flanking gene (- >1 kb; -- >10 kb; --- >100 kb).

**Table 2. Demographic characteristics of analyzed data**

		Sex (%)		Age distribution (%)			
		Males	Females	≤ 30	> 30 and ≤ 45	> 45 and ≤ 60	> 60
<b>Cilantro aversion</b>	cases	8309 (42.1%)	11422 (57.9%)	2158 (10.9%)	4645 (23.5%)	5273 (26.7%)	7655 (38.8%)
	controls	26121 (46.3%)	30239 (53.7%)	5983 (10.6%)	16325 (29.0%)	16771 (29.8%)	17281 (30.7%)
<b>Sweet preference</b>	cases	26420 (48.1%)	28481 (51.9%)	8459 (15.4%)	17181 (31.3%)	14677 (26.7%)	14584 (26.6%)
	controls	35108 (54.8%)	28941 (45.2%)	10277 (16.0%)	21054 (32.9%)	17299 (27.0%)	15419 (24.1%)
<b>Sensitivity to chewing sounds</b>	cases	7222 (41.0%)	10384 (59.0%)	3321 (18.9%)	6415 (36.4%)	4666 (26.5%)	3204 (18.2%)
	controls	31009 (49.2%)	31992 (50.8%)	5995 (9.5%)	13953 (22.1%)	18407 (29.2%)	24646 (39.1%)

Numbers (and percentages) of individuals in the analyzed data sets.

## Definitions

**Allele** – A version of a DNA sequence at a specific genomic location (either coding or non-coding). An individual's **genotype** is his/her set of alleles at a specific genomic location.

**Ancestry** – Characterization of an individual as belonging to a group with a specific historical geographic location and ethnic identity. Determined computationally by comparing the individual's genome to reference datasets for various ancestry groups.

**Effect Allele Frequency (EAF)** – The frequency in a population of the allele associated with higher odds of having a specific trait.

**Genome Wide Association Study (GWAS)** – A study that uses an array of SNPs that span the genome to identify loci linked with a particular trait.

**Hardy Weinberg Equilibrium (HWE)** – An idealized state where the allele frequencies for a specific gene in a population are constant from generation to generation. HWE may be disturbed by natural selection, genetic drift, mutation, and other factors that shift allele frequencies.

**Intron** – A region of DNA within a gene that is normally removed (spliced) after transcription and before translation to a protein.

**Linkage Disequilibrium (LD)** – A non-random association between alleles at different locations in the genome indicating shared ancestral origins and a higher likelihood of being inherited together after recombination.

**Odds Ratio (OR)** – A ratio of the odds of having a trait given one SNP genotype to the odds of having the same trait given another SNP genotype, equivalent to the effect size of a genetic association for a binary trait.

**Quantile-Quantile (Q-Q) Plot** – A graphical representation of the quantiles of one data set plotted against the quantiles of another (reference) data set. In this white paper the Q-Q plot is used to evaluate whether an experimental data distribution is significantly different from a theoretical null (no SNP-trait association) distribution.

**Regression Coefficient ( $\beta$ )** – A measure of the strength of the linear relationship between a SNP allele and a trait, used to estimate the effect size of a genetic association.

**Single Nucleotide Polymorphism (SNP)** – A single-nucleotide variation at a specific location in DNA sequence that may be used as a marker for a specific human trait or traits.

**Trait** – An easily observable characteristic that can be narrowly defined and identified in multiple individuals. In this white paper "trait" is identical to "phenotype".

## Acknowledgements

We thank the customers of 23andMe for answering surveys and participating in this research. We also thank all the employees of 23andMe, who together have made this research possible.

## References

1. Eriksson N, Macpherson JM, Tung JY, Hon LS, Naughton B, Saxonov S, et al. Web-Based, Participant-Driven Studies Yield Novel Genetic Associations for Common Traits. *PLoS Genet.* 2010;6: e1000993. doi:10.1371/journal.pgen.1000993
2. Do CB, Tung JY, Dorfman E, Kiefer AK, Drabant EM, Francke U, et al. Web-Based Genome-Wide Association Study Identifies Two Novel Loci and a Substantial Genetic Component for Parkinson's Disease. *PLoS Genet.* 2011;7: e1002141. doi:10.1371/journal.pgen.1002141
3. Tung JY, Do CB, Hinds DA, Kiefer AK, Macpherson JM, Chowdry AB, et al. Efficient Replication of over 180 Genetic Associations with Self-Reported Medical Data. *PLoS ONE.* 2011;6: e23473. doi:10.1371/journal.pone.0023473
4. Eriksson N, Benton GM, Do CB, Kiefer AK, Mountain JL, Hinds DA, et al. Genetic variants associated with breast size also influence breast cancer risk. *BMC Med Genet.* 2012;13: 53. doi:10.1186/1471-2350-13-53
5. Eriksson N, Tung JY, Kiefer AK, Hinds DA, Francke U, Mountain JL, et al. Novel Associations for Hypothyroidism Include Known Autoimmune Risk Loci. *PLoS ONE.* 2012;7: e34442. doi:10.1371/journal.pone.0034442
6. Hinds DA, McMahon G, Kiefer AK, Do CB, Eriksson N, Evans DM, et al. A genome-wide association meta-analysis of self-reported allergy identifies shared and allergy-specific susceptibility loci. *Nat Genet.* 2013;45: 907–911. doi:10.1038/ng.2686
7. Kiefer AK, Tung JY, Do CB, Hinds DA, Mountain JL, Francke U, et al. Genome-Wide Analysis Points to Roles for Extracellular Matrix Remodeling, the Visual Cycle, and Neuronal Development in Myopia. *PLoS Genet.* 2013;9: e1003299. doi:10.1371/journal.pgen.1003299
8. Nalls MA, Pankratz N, Lill CM, Do CB, Hernandez DG, Saad M, et al. Large-scale meta-analysis of genome-wide association data identifies six new risk loci for Parkinson's disease. *Nat Genet.* 2014;46: 989–993. doi:10.1038/ng.3043
9. Ferreira MAR, Matheson MC, Tang CS, Granell R, Ang W, Hui J, et al. Genome-wide association analysis identifies 11 risk variants associated with the asthma with hay fever phenotype. *J Allergy Clin Immunol.* 2014;133: 1564–1571. doi:10.1016/j.jaci.2013.10.030

10. Tung JY, Kiefer AK, Mullins M, Francke U, Eriksson N. Genome-Wide Association Analysis Implicates Elastic Microfibrils in the Development of Nonsyndromic Striae Distensae. *J Invest Dermatol.* 2013;133: 2628–2631. doi:10.1038/jid.2013.196
11. Heilmann S, Kiefer AK, Fricker N, Drichel D, Hillmer AM, Herold C, et al. Androgenetic Alopecia: Identification of Four Genetic Risk Loci and Evidence for the Contribution of WNT Signaling to Its Etiology. *J Invest Dermatol.* 2013;133: 1489–1496. doi:10.1038/jid.2013.43
12. Hromatka BS, Tung JY, Kiefer AK, Do CB, Hinds DA, Eriksson N. Genetic variants associated with motion sickness point to roles for inner ear development, neurological processes and glucose homeostasis. *Hum Mol Genet.* 2015; ddv028. doi:10.1093/hmg/ddv028
13. Chang ALS, Raber I, Xu J, Li R, Spitale R, Chen J, et al. Assessment of the Genetic Basis of Rosacea by Genome-Wide Association Study. *J Invest Dermatol.* 2015;135: 1548–1555. doi:10.1038/jid.2015.53
14. Durand EY, Do CB, Mountain JL, Macpherson JM. Ancestry Composition: A Novel, Efficient Pipeline for Ancestry Deconvolution. *bioRxiv.* 2014; 010512. doi:10.1101/010512
15. Henn BM, Hon L, Macpherson JM, Eriksson N, Saxonov S, Pe'er I, et al. Cryptic Distant Relatives Are Common in Both Isolated and Cosmopolitan Genetic Samples. *PLoS ONE.* 2012;7: e34267. doi:10.1371/journal.pone.0034267
16. Li R, Brockschmidt FF, Kiefer AK, Stefansson H, Nyholt DR, Song K, et al. Six Novel Susceptibility Loci for Early-Onset Androgenetic Alopecia and Their Unexpected Association with Common Diseases. *PLoS Genet.* 2012;8: e1002746. doi:10.1371/journal.pgen.1002746
17. Pichler I, Del Greco M. F, Gögele M, Lill CM, Bertram L, Do CB, et al. Serum Iron Levels and the Risk of Parkinson Disease: A Mendelian Randomization Study. *PLoS Med.* 2013;10: e1001462. doi:10.1371/journal.pmed.1001462
18. Rietveld CA, Conley D, Eriksson N, Esko T, Medland SE, Vinkhuyzen AAE, et al. Replicability and robustness of genome-wide-association studies for behavioral traits. *Psychol Sci.* 2014;25: 1975–1986. doi:10.1177/0956797614545132
19. Wigginton JE, Cutler DJ, Abecasis GR. A Note on Exact Tests of Hardy-Weinberg Equilibrium. *Am J Hum Genet.* 2005;76: 887–893.
20. Devlin B, Roeder K. Genomic control for association studies. *Biometrics.* 1999;55: 997–1004.

21. Zohary D, Hopf M. Domestication of Plants in the Old World: The Origin and Spread of Cultivated Plants in West Asia, Europe, and the Nile Valley. 3 edition. Oxford New York: Oxford University Press; 2001.
22. Bostock J, Riley HT. The Natural History of Pliny. BiblioLife; 2009.
23. Faas P. Around the Roman Table: Food and Feasting in Ancient Rome. 1 edition. New York, N.Y.: Palgrave Macmillan; 2002.
24. Mauer L, El-Sohehy A. Prevalence of cilantro (*Coriandrum sativum*) disliking among different ethnocultural groups. *Flavour*. 2012;1: 8. doi:10.1186/2044-7248-1-8
25. Keith R, Cadwallader, Daniel Benitez, Sompoche Pojjanapimol, Orasa Suriyaphan, Tanoj Singh. Characteristic Aroma Components of the Cilantro Mimics. *Natural Flavors and Fragrances*. American Chemical Society; 2005. pp. 117–128. Available: <http://dx.doi.org/10.1021/bk-2005-0908.ch008>
26. Eyres G, Dufour J-P, Hallifax G, Sotheeswaran S, Marriott PJ. Identification of character-impact odorants in coriander and wild coriander leaves using gas chromatography-olfactometry (GCO) and comprehensive two-dimensional gas chromatography-time-of-flight mass spectrometry (GC x GC-TOFMS). *J Sep Sci*. 2005;28: 1061–1074.
27. Quynh CTT, Kubota K. Aroma Constituents and Enzyme Activities of Japanese Long Coriander Leaves (*Culantro*, *Eryngium foetidum* L.). *Food Sci Technol Res*. 2012;18: 287–294. doi:10.3136/fstr.18.287
28. Glusman G, Bahar A, Sharon D, Pilpel Y, White J, Lancet D. The olfactory receptor gene superfamily: data mining, classification, and nomenclature. *Mamm Genome Off J Int Mamm Genome Soc*. 2000;11: 1016–1023.
29. Reed DR, Knaapila A. Genetics of Taste and Smell: Poisons and Pleasures. *Prog Mol Biol Transl Sci*. 2010;94: 213–240. doi:10.1016/B978-0-12-375003-7.00008-X
30. Kim U, Jorgenson E, Coon H, Leppert M, Risch N, Drayna D. Positional cloning of the human quantitative trait locus underlying taste sensitivity to phenylthiocarbamide. *Science*. 2003;299: 1221–1225. doi:10.1126/science.1080190
31. Keller A, Zhuang H, Chi Q, Vosshall LB, Matsunami H. Genetic variation in a human odorant receptor alters odour perception. *Nature*. 2007;449: 468–472. doi:10.1038/nature06162

32. Mainland JD, Keller A, Li YR, Zhou T, Trimmer C, Snyder LL, et al. The missense of smell: functional variability in the human odorant receptor repertoire. *Nat Neurosci.* 2014;17: 114–120. doi:10.1038/nn.3598
33. McRae JF, Jaeger SR, Bava CM, Beresford MK, Hunter D, Jia Y, et al. Identification of Regions Associated with Variation in Sensitivity to Food-Related Odors in the Human Genome. *Curr Biol.* 2013;23: 1596–1600. doi:10.1016/j.cub.2013.07.031
34. Jaeger SR, McRae JF, Bava CM, Beresford MK, Hunter D, Jia Y, et al. A Mendelian Trait for Olfactory Sensitivity Affects Odor Experience and Food Selection. *Curr Biol.* 2013;23: 1601–1605. doi:10.1016/j.cub.2013.07.030
35. Knaapila A, Hwang L-D, Lysenko A, Duke FF, Fesi B, Khoshnevisan A, et al. Genetic Analysis of Chemosensory Traits in Human Twins. *Chem Senses.* 2012; bjs070. doi:10.1093/chemse/bjs070
36. Malnic B, Godfrey PA, Buck LB. The human olfactory receptor gene family. *Proc Natl Acad Sci U S A.* 2004;101: 2584–2589. doi:10.1073/pnas.0307882100
37. Eriksson N, Wu S, Do CB, Kiefer AK, Tung JY, Mountain JL, et al. A genetic variant near olfactory receptor genes influences cilantro preference. *Flavour.* 2012;1: 22. doi:10.1186/2044-7248-1-22
38. Kurland MD, Newcomer MB, Peterlin Z, Ryan K, Firestein S, Batista VS. Discrimination of Saturated Aldehydes by the Rat I7 Olfactory Receptor. *Biochemistry (Mosc).* 2010;49: 6302–6304. doi:10.1021/bi100976w
39. Araneda RC, Kini AD, Firestein S. The molecular receptive range of an odorant receptor. *Nat Neurosci.* 2000;3: 1248–1255. doi:10.1038/81774
40. Araneda RC, Peterlin Z, Zhang X, Chesler A, Firestein S. A pharmacological profile of the aldehyde receptor repertoire in rat olfactory epithelium. *J Physiol.* 2004;555: 743–756. doi:10.1113/jphysiol.2003.058040
41. Krautwurst D, Yau KW, Reed RR. Identification of ligands for olfactory receptors by functional expression of a receptor library. *Cell.* 1998;95: 917–926.
42. Li Y, Peterlin Z, Ho J, Yarnitzky T, Liu MT, Fichman M, et al. Aldehyde Recognition and Discrimination by Mammalian Odorant Receptors via Functional Group-Specific Hydration Chemistry. *ACS Chem Biol.* 2014; doi:10.1021/cb400290u
43. Li X, Staszewski L, Xu H, Durick K, Zoller M, Adler E. Human receptors for sweet and umami taste. *Proc Natl Acad Sci.* 2002;99: 4692–4696. doi:10.1073/pnas.072090199

44. Chandrashekar J, Hoon MA, Ryba NJP, Zuker CS. The receptors and cells for mammalian taste. *Nature*. 2006;444: 288–294. doi:10.1038/nature05401
45. Garcia-Bailo B, Toguri C, Eny KM, El-Sohemy A. Genetic variation in taste and its influence on food selection. *Omics J Integr Biol*. 2009;13: 69–80. doi:10.1089/omi.2008.0031
46. Keskitalo K, Tuorila H, Spector TD, Cherkas LF, Knaapila A, Silventoinen K, et al. Same genetic components underlie different measures of sweet taste preference. *Am J Clin Nutr*. 2007;86: 1663–1669.
47. Keskitalo K, Knaapila A, Kallela M, Palotie A, Wessman M, Sammalisto S, et al. Sweet taste preferences are partly genetically determined: identification of a trait locus on chromosome 16. *Am J Clin Nutr*. 2007;86: 55–63.
48. Beauchamp GK, Moran M. Dietary experience and sweet taste preference in human infants. *Appetite*. 1982;3: 139–152. doi:10.1016/S0195-6663(82)80007-X
49. Beauchamp GK, Moran M. Acceptance of sweet and salty tastes in 2-year-old children. *Appetite*. 1984;5: 291–305. doi:10.1016/S0195-6663(84)80002-1
50. Bertino M, Beauchamp GK, Engelman K. Increasing dietary salt alters salt taste preference. *Physiol Behav*. 1986;38: 203–213. doi:10.1016/0031-9384(86)90155-1
51. Bertino M, Beauchamp GK, Jen KC. Rated taste perception in two cultural groups. *Chem Senses*. 1983;8: 3–15. doi:10.1093/chemse/8.1.3
52. Prescott J. Comparisons of taste perceptions and preferences of Japanese and Australian consumers: overview and implications for cross-cultural sensory research. *Food Qual Prefer*. 1998;9: 393–402. doi:10.1016/S0950-3293(98)00021-4
53. Lanfer A, Bammann K, Knof K, Buchecker K, Russo P, Veidebaum T, et al. Predictors and correlates of taste preferences in European children: The IDEFICS study. *Food Quality and Preference*. Food Quality and Preference; 2013. Available: <http://dx.doi.org/10.1016/j.foodqual.2012.09.006>
54. Sclafani A. Carbohydrate taste, appetite, and obesity: An overview. *Neurosci Biobehav Rev*. 1987;11: 131–153. doi:10.1016/S0149-7634(87)80019-2
55. Mattes RD. Fat taste and lipid metabolism in humans. *Physiol Behav*. 2005;86: 691–697. doi:10.1016/j.physbeh.2005.08.058
56. Margolskee RF. Molecular Mechanisms of Bitter and Sweet Taste Transduction. *J Biol Chem*. 2002;277: 1–4. doi:10.1074/jbc.R100054200



57. Mace OJ, Affleck J, Patel N, Kellett GL. Sweet taste receptors in rat small intestine stimulate glucose absorption through apical GLUT2. *J Physiol.* 2007;582: 379–392. doi:10.1113/jphysiol.2007.130906
58. Eny KM, Wolever TMS, Fontaine-Bisson B, El-Sohemy A. Genetic variant in the glucose transporter type 2 is associated with higher intakes of sugars in two distinct populations. *Physiol Genomics.* 2008;33: 355–360. doi:10.1152/physiolgenomics.00148.2007
59. Kharitonov A, Shiyanova TL, Koester A, Ford AM, Micanovic R, Galbreath EJ, et al. FGF-21 as a novel metabolic regulator. *J Clin Invest.* 2005;115: 1627–1635. doi:10.1172/JCI23606
60. Fisher M, Ffolliott M, Chui PC, Antonellis PJ, Bina HA, Kharitonov A, Flier JS, et al. Obesity Is a Fibroblast Growth Factor 21 (FGF21)-Resistant State. *Diabetes.* 2010;59: 2781–2789. doi:10.2337/db10-0193
61. Woo YC, Xu A, Wang Y, Lam KSL. Fibroblast Growth Factor 21 as an emerging metabolic regulator: clinical perspectives. *Clin Endocrinol (Oxf).* 2013;78: 489–496. doi:10.1111/cen.12095
62. Emanuelli B, Vienberg SG, Smyth G, Cheng C, Stanford KI, Arumugam M, et al. Interplay between FGF21 and insulin action in the liver regulates metabolism. *J Clin Invest.* 2014;124: 515–527. doi:10.1172/JCI67353
63. Zhang X, Yeung DCY, Karpisek M, Stejskal D, Zhou Z-G, Liu F, et al. Serum FGF21 Levels Are Increased in Obesity and Are Independently Associated With the Metabolic Syndrome in Humans. *Diabetes.* 2008;57: 1246–1253. doi:10.2337/db07-1476
64. Chen C, Cheung BM, Tso AWK, Wang Y, Law LSC, Ong KL, et al. High Plasma Level of Fibroblast Growth Factor 21 Is an Independent Predictor of Type 2 Diabetes A 5.4-year population-based prospective study in Chinese subjects. *Diabetes Care.* 2011;34: 2113–2115. doi:10.2337/dc11-0294
65. Zhao Y, Dunbar JD, Kharitonov A. FGF21 as a Therapeutic Reagent. In: MD MK, editor. *Endocrine FGFs and Klothos.* Springer US; 2012. pp. 214–228. Available: [http://link.springer.com/chapter/10.1007/978-1-4614-0887-1\\_14](http://link.springer.com/chapter/10.1007/978-1-4614-0887-1_14)
66. Iglesias P, Selgas R, Romero S, Díez JJ. MECHANISMS IN ENDOCRINOLOGY: Biological role, clinical significance, and therapeutic possibilities of the recently discovered metabolic hormone fibroblastic growth factor 21. *Eur J Endocrinol.* 2012;167: 301–309. doi:10.1530/EJE-12-0357
67. Fawcett KA, Barroso I. The genetics of obesity: FTO leads the way. *Trends Genet.* 2010;26: 266–274. doi:10.1016/j.tig.2010.02.006

68. Hess ME, Brüning JC. The fat mass and obesity-associated (FTO) gene: Obesity and beyond? *Biochim Biophys Acta BBA - Mol Basis Dis.* 2014;1842: 2039–2047. doi:10.1016/j.bbadis.2014.01.017
69. Smemo S, Tena JJ, Kim K-H, Gamazon ER, Sakabe NJ, Gómez-Marín C, et al. Obesity-associated variants within FTO form long-range functional connections with IRX3. *Nature.* 2014;507: 371–375. doi:10.1038/nature13138
70. Ragvin A, Moro E, Fredman D, Navratilova P, Drivenes Ø, Engström PG, et al. Long-range gene regulation links genomic type 2 diabetes and obesity risk regions to HHEX, SOX4, and IRX3. *Proc Natl Acad Sci.* 2010;107: 775–780. doi:10.1073/pnas.0911591107
71. Chu AY, Workalemahu T, Paynter NP, Rose LM, Giulianini F, Tanaka T, et al. Novel locus including FGF21 is associated with dietary macronutrient intake. *Hum Mol Genet.* 2013; ddt032. doi:10.1093/hmg/ddt032
72. Jastreboff PJ, Jastreboff MM. Tinnitus retraining therapy for patients with tinnitus and decreased sound tolerance. *Otolaryngol Clin North Am.* 2003;36: 321–336.
73. Schröder A, Vulink N, Denys D. Misophonia: Diagnostic Criteria for a New Psychiatric Disorder. *PLoS ONE.* 2013;8: e54706. doi:10.1371/journal.pone.0054706
74. Baguley DM. Hyperacusis. *J R Soc Med.* 2003;96: 582–585.
75. Wu MS, Lewin AB, Murphy TK, Storch EA. Misophonia: Incidence, Phenomenology, and Clinical Correlates in an Undergraduate Student Sample. *J Clin Psychol.* 2014;70: 994–1007. doi:10.1002/jclp.22098
76. Rubin BP, Tucker RP, Martin D, Chiquet-Ehrismann R. Teneurins: A Novel Family of Neuronal Cell Surface Proteins in Vertebrates, Homologous to the *Drosophila* Pair-Rule Gene Product Ten-m. *Dev Biol.* 1999;216: 195–209. doi:10.1006/dbio.1999.9503
77. Silva J-P, Lelianova VG, Ermolyuk YS, Vysokov N, Hitchen PG, Berninghausen O, et al. Latrophilin 1 and its endogenous ligand Lasso/teneurin-2 form a high-affinity transsynaptic receptor pair with signaling capabilities. *Proc Natl Acad Sci.* 2011;108: 12113–12118. doi:10.1073/pnas.1019434108
78. Mosca TJ, Hong W, Dani VS, Favaloro V, Luo L. Trans-synaptic Teneurin signalling in neuromuscular synapse organization and target choice. *Nature.* 2012;484: 237–241. doi:10.1038/nature10923
79. Jastreboff PJ. Tinnitus Retraining Therapy. In: Møller AR, Langguth B, Ridder DD, Kleinjung T, editors. *Textbook of Tinnitus.* Springer New York; 2011. pp.

575–596. Available: [http://link.springer.com/chapter/10.1007/978-1-60761-145-5\\_73](http://link.springer.com/chapter/10.1007/978-1-60761-145-5_73)

## Revision History

Originally published on December 4, 2014.

Revision A published February 25, 2015. The following changes were made in Revision A:

- Added new GWAS data for sweet taste preference to Table 1 and new Figure 2
- Corrected EAF values for cilantro preference SNPs in Table 1, which were listed for the wrong allele
- Added Table 2 – Demographic characteristics

Revision B published June 3, 2015. The following changes were made in Revision B:

- Added new GWAS data for chewing sound sensitivity to Table 1 and Table 2, and new Figure 3
- Added recent 23andMe publications to references